

Name: _____

Period: _____

Momentum In Class Review


- A. $p_{1+2B} = p_{1A} + p_{2A}$
- B. $p_B - I = 0$
- C. $0 = p_{1A} + p_{2A}$
- D. $p_B + I = p_A$
- E. $p_{1B} + p_{2B} = p_{1A} + p_{2A}$
- F. $p_{1B} + p_{2B} = p_{1+2A}$

- 1. A car speeds up.
- 2. A person running catches a football.
- 3. Two moving cars hit and bounce off.
- 4. A moving airplane drops a bomb.
- 5. A rocket at rest turns on its engine: hot gases go back; the rocket goes forward.
- 6. A moving car uses its brakes to stop.

- 7. Which has more momentum?
 - A. A fast baseball or a slow baseball?
 - B. A bowling ball or a baseball with the same speed?
 - C. A slow ping pong ball or a house?
- 8. Give two ways momentum can change.

9. Does a large force always cause a large impulse? Explain.

10. 15 N acts for 8 seconds. How much momentum was gained?




- 11. How much momentum was gained above?
- 12. How big is the impulse acting on the object?
- 13. Calculate the time the force acted.
- 14. Calculate the acceleration of the object.
- 15. What is the final velocity of the object?

16. Elastic, Inelastic, or Perfectly Inelastic (could be more than one)?

- A. _____ $\Sigma p_{\text{before}} = \Sigma p_{\text{after}}, \Sigma E_{k\text{before}} \neq \Sigma E_{k\text{after}}$
- B. _____ $\Sigma p_{\text{before}} = \Sigma p_{\text{after}}, \Sigma E_{k\text{before}} = \Sigma E_{k\text{after}}$
- C. _____ $\Sigma p_{\text{before}} = \Sigma p_{\text{after}}, \text{ and } m_{\text{after}} = m_{1+2}$
- D. _____ There is little or no sound.
- E. _____ There is a lot of noise.
- F. _____ The objects are mangled, or crushed.




- 17. Two objects collide as shown above. They don't stick.
 - A. What happens to the momentum of the 4 kg object?
 - B. What happens to the momentum of the 6 kg object?
 - C. What happens to the total momentum of the system?



- 18. Two identical carts moving 6 m/s stop. The Cart 1 hits a spring. The Cart 2 just hits a wall.
 - A. Calculate the initial momentum of the carts.
 - B. Calculate the change of momentum of the carts.
 - C. Which cart experienced the bigger change of momentum?
 - D. Which cart felt the bigger impulse? E. Which cart felt the bigger force?
 - F. Calculate the force on each cart.
 - G. So, to give the same Δp you have two choices:

Name: _____

Period: _____




19. Slim Jim is also an astronaut. His space ship "Galactic Cruiser" is at rest when he shoots his space cannon.

- A. What is the mass of the ship?
- B. What is the weight of the ship?
- C. Calculate the final velocity of the ship.


D. Which has more momentum afterwards: the ship or the projectile?

20. A 20 g bullet is shot 800 m/s into 50 kg object. What is the final speed of the combined object?


- A. If $1000\text{ g} = 1\text{ kg}$, what is the mass of the bullet in kilograms?
- B. What is the mass of the *combined object*?
- C. Under the diagram, calculate the final speed.
- D. The numbers given are realistic for a bullet and a person. In movies, a bullet causes a person to be thrown backwards violently. How likely is the movie scenario? Explain.



Graph 1 Force vs. Time



Graph 2 Force vs. Time



21. Use the graphs above to answer the following questions.

22. Graph 1 or Graph 2?

- A. ____ Shows an object with a positive acceleration
- B. ____ Could be an object moving to the right and slowing down.
- C. ____ Shows a negative change of speed.
- D. ____ Shows a force pushing to the left.

24. Find the impulse of Graph 1.

23. Force A, B, C, D, or E (could be more than one)?

- A. ____ Is the strongest positive force.
- B. ____ Is the greatest negative force.
- C. ____ Is the weakest positive force.
- D. ____ Will cause the fastest negative acceleration.
- E. ____ Is the strongest force pulling left.
- F. ____ Shows negative acceleration.

25. If a 2 kg object going 6 m/s feels the impulse on Graph 1, find its final velocity.

Momentum In Class Review

- A. $p_{1+2B} = p_{1A} + p_{2A}$ 4
 B. $p_B - I = 0$ 6
 C. $0 = p_{1A} + p_{2A}$ 5
 D. $p_B + I = p_A$ 1
 E. $p_{1B} + p_{2B} = p_{1A} + p_{2A}$ 3
 F. $p_{1B} + p_{2B} = p_{1+2A}$ 2

1. D A car speeds up.
 2. F A person running catches a football.
 3. E Two moving cars hit and bounce off.
 4. A A moving airplane drops a bomb.
 5. C A rocket at rest turns on its engine: hot gases go back; the rocket goes forward.
 6. B A moving car uses its brakes to stop.

7. Which has more momentum?
 A. A fast baseball or a slow baseball?
 B. A bowling ball or a baseball going 2 m/s?
 C. A slow ping pong ball or a house?
 8. Give two ways momentum can change.
 transferred (collision)
 Impulse (force)


9. Does a large force always cause a large impulse? Explain.

No, if the time is small.

10. 15 N acts for 8 seconds. How much momentum was gained?

$$I = Ft = \Delta p$$

$$15(8) = 120 \text{ kgm/s}$$



11. How much momentum was gained above?

$$36 - 12 = 24 \text{ kgm/s}$$

12. How big is the impulse acting on the object? 24 kgm/s

13. Calculate the time the force acted.

$$24 = Ft$$

$$24 = 8t \quad t = 3 \text{ sec}$$

14. Calculate the acceleration of the object.

$$F = ma$$

$$8 = 4a \quad a = 2 \text{ m/s}^2$$

$$a = \frac{\Delta v}{t} = \frac{9-3}{3} = \frac{6}{3} = 2 \text{ m/s}^2$$

15. What is the final velocity of the object?

$$\frac{36}{4} = 9 \text{ m/s} \quad p = mv \text{ so } v = p/m$$


16. Elastic, Inelastic, or Perfectly Inelastic (could be more than one)?

- A. I, PI $\Sigma p_{\text{before}} = \Sigma p_{\text{after}}, \Sigma E_{k\text{before}} \neq \Sigma E_{k\text{after}}$
 B. E $\Sigma p_{\text{before}} = \Sigma p_{\text{after}}, \Sigma E_{k\text{before}} = \Sigma E_{k\text{after}}$
 C. PI $\Sigma p_{\text{before}} = \Sigma p_{\text{after}}, \text{ and } m_{\text{after}} = m_{1+2}$ ← they stick
 D. E There is little or no sound.
 E. I, PI There is a lot of noise.
 F. I, PI The objects are mangled, or crushed.



17. Two objects collide as shown above.

- A. What happens to the momentum of the 4 kg object?
gained (increases)
 B. What happens to the momentum of the 6 kg object?
loses
 C. What happens to the total momentum of the system?
stays same



18. Two identical carts moving 6 m/s stop. The left cart hits a spring. The right cart just hits a wall.

A. Calculate the initial momentum of the carts. 9 kgm/s

B. Calculate the change of momentum of the carts. -9 kgm/s (stop)

C. Which cart experienced the bigger change of momentum? same

D. Which cart felt the bigger impulse? same

E. Which cart felt the bigger force? cart 2


F. Calculate the force on each cart.

$$I = Ft \quad F = -18 \text{ N} \quad F = -9(0.1) = -90 \text{ N}$$

$$-9 = F(0.5)$$

G. So, to give the same Δp you have two choices:


Big F, small t and vice versa





19. Slim Jim is also an astronaut. His space ship "Galactic Cruiser" is at rest when he shoots his space cannon.
- What is the mass of the ship? $2,500 \text{ kg}$
 - What is the weight of the ship? 0 N

- C. Calculate the final velocity of the ship.
- $$0 = p_1 + p_2$$
- $$0 = 2,500v + 1.5(800)$$
- $$-1200 = 2,500v \quad v = \frac{12}{25} = -4 \text{ m/s}$$
- D. Which has more momentum afterwards: the ship or the projectile? *same (just opp.)*

20. A 20 g bullet is shot 800 m/s into 50 kg object. What is the final speed of the combined object?
- If $1000 \text{ g} = 1 \text{ kg}$, what is the mass of the bullet in kilograms? 0.02 kg
 - What is the mass of the **combined object**? 50.02 kg
 - Under the diagram, calculate the final speed.
 - The numbers given are realistic for a bullet and a person. In movies, a bullet causes a person to be thrown backwards violently. How likely is the movie scenario? Explain.



Not possible - bullet is so light that it can't give enough p.



21. Use the graphs above to answer the following questions.

22. Which Graph?
- 1 Shows an object with a positive acceleration
 - 2 Could be an object moving to the right and slowing down. *Fisney?*
 - 2 Shows a negative change of speed.
 - 2 Shows a force pushing to the left.

23. Which force (or forces)?

- B Is the strongest positive force.
- E Is the greatest negative force.
- C Is the weakest positive force.
- E Will cause the fastest negative acceleration.
- E Is the strongest force pulling left.
- D, E Shows negative acceleration.

24. Find the impulse of Graph 1.

$$= \frac{1}{2}(1)(5) + 3(5) + \frac{1}{2}(5)(0.5)$$

$$= 2.5 + 15 + 1.25$$

$$= 18.75 \text{ kg m/s}$$

25. If a 2 kg object going 6 m/s feels the impulse on Graph 1, find its final velocity.

$$p_B + I = p_A$$

$$2(6) + 18.75 = 2(v)$$

$$30.75 = 2v$$

$$v = 15.375 \text{ m/s}$$